



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 12 AUG 2004

WIPO


PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated

2 August 2004

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

The
Patent
Office

1/77
20 JUN 03 E816461-81002732
P01/7700 0.00-031341.9

THE PATENT OFFICE
M

Request for grant of a patent

19 JUN 2003

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

LONDON

The Patent Office

Cardiff Road
Newport
South Wales
NP9 1RH

1. Your reference

A1528

2. Patent application number

(The Patent Office will fill in this part)

0314341.9

19 JUN 2003

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

ABERDEEN UNIVERSITY
RESEARCH AND INNOVATION
UNIVERSITY OFFICE
KINGS COLLEGE
ABERDEEN AB24 3FX
UK

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the country/state of its incorporation

8642183001

4. Title of the invention

A METHOD FOR PRODUCING A DI-SUBSTITUTED
PYRIDINIUM COMPOUND

5. Name of your agent (*if you have one*)

"Address for service" in the United Kingdom
to which all correspondence should be sent
(*including the postcode*)

ABLETT & STEBBING
CAPARO HOUSE
101-103 BAKER STREET
LONDON
W1U 6FQ

Patents ADP number (*if you know it*)

00006551001 ✓

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number
(*if you know it*)

Date of filing
(*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if*

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES

Patents Form 1/77

Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form	0
Description	15 ✓
Claim(s)	5 ✓ W~
Abstract	0
Drawing(s)	0

10. If you are also filing any of the following, state how many against each item.

Priority documents	0
Translations of priority documents	0
Statement of inventorship and right to grant of a patent (Patents Form 7/77)	0
Request for preliminary examination and search (Patents Form 9/77)	0
Request for substantive examination (Patents Form 10/77)	0
Any other documents (please specify)	0

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

ABLETT & STEBBING

19 June 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

G K ABLETT / S J SUER

020-7935-7720

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

A METHOD FOR PRODUCING A DI-SUBSTITUTED PYRIDINIUM COMPOUND

The present invention relates to a method for producing a di-substituted pyridinium compound.

5

Dialkylpyridinium compounds, (Di-APS), in particular 1,3-APS oligomers, are known to be produced by sponges, for example the Haploscerid genera such as *Haliclona*, *Amphimedon* and *Callyspongia*, as part of their chemical defences, and have 10 potentially useful biological properties. Diverse biological activities have been identified for different 1,3-APS compositions, including cytotoxicity, neurotoxicity and inhibition of action potentials, stimulation of transmitter release, inhibition of K⁺ conductances, and anticholinesterase 15 activity. At least some of these observed actions of 1,3-APS compositions relate to the pore forming or membrane lesion effects of these compounds, which properties may be useful, for example, in the transfection of cells with genetic material.

20

Naturally occurring 1,3-APS compounds are produced as cocktails of different 1,3-APS compounds, from which individual 1,3-APS compounds are difficult to isolate, due to the same basic structure and very similar molecular weights 25 of the different 1,3-APS compounds. However, it is desirable to be able to isolate different individual 1,3-APS compounds, in order to determine their different biological activities, for example with regard to the effect of degree of polymerisation, and length and rigidity of linking chains.

30

An alternative approach to isolating individual naturally occurring Di-APS compounds, is to synthesise these compounds by laboratory methods. Di-APS compounds generally occur as

high molecular weight linear oligomers, having a molecular weight ranging from 1 KDa to greater than 25 KDa, having varying lengths of aliphatic chains linking the pyridine units. However, model linear Di-APS compounds closely
5 analogous to the natural products remain to be synthesized by laboratory methods in a controllable fashion.

Previous studies have succeeded in producing linear 1,3-APS oligomers, but the methods employed meant that the product
10 obtained consisted of a mixture of linear and cyclic oligomers with a wide range of molecular weights (see for example, Davies-Coleman, M.T.; Faulkner, D.J. *J. Org. Chem.* 1993, 58, 5925-5930). The approach used in these studies was to synthesize the 3-substituted pyridine monomer and introduce
15 a good leaving group, for example bromide, at the end of the alkyl chain. This monomer was then refluxed to initiate polymerisation. Alternative methods based on this strategy incorporate an ether linkage in the linking chain (Gil, L. et al *Tetrahedron Lett.* 1995, 36, 2059-2062), or form cyclic
20 dimers (Morimoto, Y.; Yokoe, C. *Tetrahedron Lett.* 1997, 38, 8981-8984) or tetramers with short linking chains (Shinoda, S. et al, *Chem. Commun.* 1998, 181).

A further study for producing macrocyclic 1,3-APS compounds
25 is disclosed in Kaiser, A. et al., *J. Am. Chem. Soc.* 1998, 120, 8026-8034, which involves the formation and subsequent reaction of an N-(2,4-dinitrobenzene) pyridinium salt (Zinke salt) to yield linear 1,3-APS compounds. However, this method has certain disadvantages, in that it is relatively complex,
30 and has only been shown to work for small macrocyclic oligomers.

Accordingly, there still exists a need for a reliable method

for synthesizing linear di-substituted pyridinium compounds, for example di-APS oligomers and polymers. The present invention seeks to provide such a method, which overcomes the aforementioned disadvantages of previous methods.

5

According to the present invention there is provided a method of producing a linear di-substituted pyridinium compound, the method comprising the steps of:

(a) attaching a first 2, 3, or 4-substituted pyridine
10 compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-X}$ (1) to a solid support, to form a compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-Y-SUPPORT}$ (2), wherein SUPPORT represents the solid support, R is selected from hydrogen, hydroxyl, and substituted or unsubstituted alkyl, alkoxy, aryl, alkaryl, aralkyl, and alkenyl groups, R' is a
15 first linking group, X is a group which can react with the solid support to attach the first pyridine compound to the support, and Y is absent or is a second linking group;

(b) forming a di-substituted pyridine compound of the formula $\text{A-}^+\text{NC}_5\text{R}_4\text{-R}'\text{-Z}$ (3) from a second 2, 3, or 4-substituted
20 pyridine compound of formula (1), which second pyridine compound may be the same as or different to the first pyridine compound, wherein A is a protecting group, and Z is a leaving group;

(c) reacting the compound of formula (2) formed in step
25 (a) with the compound of formula (3) formed in step (b), to form a di-substituted pyridinium compound of the formula $\text{A-}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}']_n\text{Y-SUPPORT}$ (4), wherein Q⁺ is a counter ion and n=1;

(d) optionally, repeating step (c) as many times as
30 required to obtain a compound of formula (4) wherein n is an integer of 2 or greater; and

(e) detaching the compound of formula (4) from the solid support, and removing the protecting group A to form a

di-substituted pyridinium compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{-X}$ (5), wherein n is an integer, and Q^- and X are a counter ion and a group which can react with the solid support to attach the first pyridine compound to the support respectively, which may be the same or different as Q and X defined above in steps (c) and (a) respectively.

Thus, during steps (a) to (d) of the method of the present invention, one end of the compound which is to become the product compound of formula (5) is bound to a solid support, thereby removing it from the reaction equilibrium, and substantially eliminating cyclisation. The use of protecting and leaving groups per the method of the present invention allows for the controlled addition of monomer units without macrocycle formation occurring in solution. The method of the present invention can be used to produce di-substituted pyridinium compounds in high rates of conversion, and in high yields. The method of the present invention also has the advantage that a variety of linking groups may be used, which may incorporate different functionalities, for example functionalities which can restrict the conformational space accessible by the di-substituted pyridinium compound, such as a double bond or cyclopropyl ring, or groups which allow the attachment of fluorescent labels, so that the di-substituted pyridinium compound can be localised intracellularly and in cell membranes.

Herein the term "di-substituted pyridinium compound" can refer to either a 1,2-, 1,3-, or 1,4-substituted pyridinium compound. However, the preferred products of the method of the present invention are 1,3-substituted pyridinium compounds, in particular linked 1,3-dialkyl pyridinium compounds.

Although the method of the present invention is suitable for producing oligomers and polymers potentially having a wide range of degrees of polymerisation, and hence molecular weights, for example $n=1$ to 20 in formula (5) above, it is particularly useful for producing oligomers having higher degrees of polymerisation, for example $n=20$ to 100 in formula (5) above.

Referring to the steps of the method of the present invention in turn, in step (a) a first 2,3, or 4-substituted, preferably 3-substituted, pyridine compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-X}$ (1) is attached to a solid support, to form a compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-Y-SUPPORT}$ (2), wherein SUPPORT represents the solid support, R is selected from hydrogen, and substituted or unsubstituted alkyl, aryl, alkaryl, aralkyl, and alkenyl groups, R' is a first linking group, X is a group which can react with the solid support to attach the first pyridine compound to the support, and Y is absent or is a second linking group. Each R group is preferably a hydrogen atom.

20

Compounds of formula (1) are commercially available, or may be synthesised as is known in the art. Thus, for example, a compound of formula (1) may be synthesised by reaction of a compound of formula $\text{Z}'\text{-R}''\text{-X}$ with a suitable pyridine compound, with protection of the X- group as necessary, wherein R'' is a linking group and Z' is a suitable leaving group. For example, a compound of formula (1) may be prepared by reacting $\text{Br-R}''\text{-OH}$ (i.e. $\text{Z}'=\text{Br}$ and $\text{X}=\text{OH}$) with t-butyldimethylchlorosilane (TBDMSCl) to form $\text{Br-R}''\text{-OTBDMS}$, which may be reacted with 2,3, or 4-methylpyridine (2,3, or 4-picoline) with deprotection of the X group to form $\text{NC}_5\text{H}_4\text{-R-OH}$, wherein R' is as defined above, i.e. a linking group equivalent to R'' plus one carbon atom. Of course, protecting groups other than

TBDMSCl may be used, as will be apparent to those skilled in the art.

As referred to above, the method of the present invention is advantageous in that a variety of linking groups may be used, which may incorporate different functionalities. Preferred R' groups have no terminal carbon atoms, which further restricts the possibility of cyclisation occurring.

Thus, in formulas (1) to (5) above, each group R' may be the same or different and selected from an alkylene group (for example, a group $-(CH_2)_m-$, wherein m is an integer from 2 to 12, preferably from 6 to 10), an alkenyl-containing group (for example, a group having from 2 to 12 carbon atoms, preferably from 6 to 10 carbon atoms, containing one or more alkenyl groups, e.g. cis- or trans- 2-butenyl, 3-hexenyl, 2,5-heptendi-yl, and 4-octenyl groups), a cyclopropanyl-containing group (for example, cis- or trans- $-(CH_2)_p$ -cyclopropanyl- $(CH_2)_q-$ wherein p and q are the same or different and are integers from 1 to 4, preferably 1 or 2). As discussed above, R' may also be a group which comprises a fluorescent label, so that the di-substituted pyridinium compound can be localised intracellularly and in cell membranes (for example, a linking group R' having a pendant alcohol group, for attachment of a fluorescent group).

Where appropriate, cis- or trans- isomerism may be introduced into linking group R' as is known in the art (for example, Morimoto, Y.; Yokoe, C. *Tetrahedron Lett.* 1997, 38, 8981-8984). A double bond may be converted into a cyclopropyl ring using carbene chemistry.

In the compound of formula (1), X is a group which can react

with the solid support to attach the compound of formula (1) to the support. Thus, the particular group X to be used in a particular compound of formula (1) will depend upon the particular solid support being used. Thus, suitable X groups
5 include hydroxyl, carboxyl, thiol, and amine groups. However, X is preferably a hydroxyl group, and particularly preferred compounds of formula (1) are pyridines 3-substituted by a hydroxyalkyl group, i.e. compounds of the formula $\text{NC}_5\text{R}_4-(\text{CH}_2)_n-\text{OH}$, wherein n is as defined above.

10

The solid support used in the present invention may be any suitable support to which the compound of formula (1) may be attached. Preferred solid support materials are organic resins having functionality which can react with group X of the
15 compound of formula (1), described above. Particularly preferred solid support materials are trityl chloride and functionalised polystyrene resins (for example, Merryfield resins, i.e. chloromethylstyrene-divinylbenzene resins).

20 Group Y in formula (2) is absent or a second linking group, depending on the particular reaction which occurs between the compound of formula (1) and the solid support to form the compound of formula (2). Thus, group X in formula (1) may be eliminated, or, for example, in the case of a compound of
25 formula (1) in which X is a hydroxyl group, the compound of formula (2) may have the more specific formula $\text{NC}_5\text{R}_4-\text{R}'-\text{O}-\text{SUPPORT}$, i.e. Y may be an oxygen atom.

In step (b) of the method of the present invention, a
30 disubstituted pyridine compound of the formula $\text{A}^+-\text{NC}_5\text{R}_4-\text{R}'-\text{Z}$ (3) is formed from a second 2,3 or 4-substituted pyridine compound of formula (1), which second pyridine compound may be the same as or different to the first pyridine compound,

wherein A is a protecting group, and Z is a leaving group. Step (b) is independent of step (a), and accordingly may be performed before, after, or simultaneously with step (a).

5 Step (b) itself thus includes two steps: a first step of converting group X to a leaving group Z, for reaction with the nitrogen atom of the pyridine group of the compound of formula (2), and a second step of protecting the nitrogen atom of the second pyridine compound, with a protecting group A.

10

In step (b), group X may be converted to a suitable leaving group Z as is known in the art. For example, as a hydroxyl group, X may be converted to a mesyl (methanesulphonyl) group by reaction with mesyl chloride. Other suitable leaving groups
15 will be apparent to those skilled in the art.

The nitrogen atom of the second pyridine compound of formula (1) used in step (b) may be protected by conversion to the N-oxide as is known in the art, i.e. protecting group A is
20 oxygen, for example by reaction of the nitrogen atom of the pyridine group with a peracid, for example m-chloroperbenzoic acid. Alternatively, the nitrogen atom may be protected by formation of the borane, i.e. A is BH_3^- .

25 In step (c) of the method of the present invention, the compound of formula (2) formed in step (a) is reacted with the compound of formula (3) formed in step (b), to form a di-substituted pyridinium compound of the formula $A^+NC_5R_4-R'-[Q^+NC_5R_4-R']_nY-SUPPORT$ (4), wherein Q^- is a counter
30 ion and $n=1$. Thus, in step (c) two pyridine-containing monomer units are combined to form a dimeric di-substituted pyridinium compound, the leaving group Z of the compound of formula (3) reacting with the nitrogen atom of the pyridine group of the

compound of formula (2). The reaction may take place on heating, in the presence of a suitable counter ion Q^- , for example a chloride, iodide, or other suitable counter ion.

5 In step (d) of the method of the present invention, step (c) is optionally repeated as many times as required to obtain a compound of formula (4) wherein n is an integer of 2 or greater. If the desired di-substituted pyridinium compound end product is a dimer (i.e. a compound of formula (5) in which
10 $n=1$), then no repetitions of step (c) are required. However, as discussed above, the method of the present invention is particularly useful for producing oligomers having higher degrees of polymerisation, for example $n=20$ to 100 in formula (5) above, which will of course require the repetition of step
15 (c) the appropriate number of times.

In preferred embodiments of the present invention, chain extension to produce oligomers having a higher degree of polymerisation may be accelerated by releasing oligomers
20 having a lower degree of polymerisation from the solid support as compounds having the formula (5), and reintroducing them as reagents as an alternative to, or in addition to, the second pyridine compound used in step (b). Thus, in these preferred embodiments, a compound of formula
25 $NC_5R_4-R'-[Q^+NC_5R_4-R']_n-X$ (5) may be converted to a compound of formula $A^--^+NC_5R_4-R'-[Q^+NC_5R_4-R']_n-Z$ (5a) per step (b), and the compound of formula (5a) may then be reacted with the compound of formula (2) formed in step (a) or (c), per step (d).

30 In step (e) of the method of the present invention, the compound of formula (4) is detached from the solid support, and reduced to form a di-substituted pyridinium compound of the formula $NC_5R_4-R'-[Q^+NC_5R_4-R']_n-X$ (5), wherein n is an


integer, and Q^- and X are a counter ion and a group which can react with the solid support to attach the first pyridine compound to the support respectively. Q^- and X of step (e) may be the same or different as Q^- and X defined above with
5 reference steps (a) and (c) respectively. The compound of formula (4) may be detached from the solid support according to the particular solid support being used. Thus, in the case of a functionalised trityl chloride resin solid support, the compound of formula (4) may be detached using a strong acid.
10 For example, if the strong acid used to detach the formula (4) from the solid support is hydrochloric acid, then counter ion Q^- will be chloride. The protecting group A can be removed as is known in the art, for example by reducing the N-oxide to a nitrogen atom where A in formula (4) is oxygen.

15

An embodiment of the present invention will now be described in detail.

Reaction Scheme 1 below describes a preferred method for
20 producing a preferred linear 1,3-alkylpyridinium compound of the present invention.

Scheme 1



wherein: Ms=methanesulphonyl (mesyl)

Et=ethyl

MCPBA = m-chloroperbenzoic acid

5 Referring to Scheme 1, 3-substituted pyridine compound (1) is formed by firstly reacting Br-R''-OH with t-butyldimethylchlorosilane (TBDMSCl) to form Br-R''-OTBDMS, wherein R'' is a suitable linking group (described hereinabove). The Br-R''-OTBDMS is then reacted with 3-methylpyridine
10 (3-picoline) with deprotection of the X group to form NC₅R₄-R'-OH (3-substituted pyridine compound (1)), wherein R' is a linking group equivalent to R'' plus one carbon atom.

A first molecule of 3-substituted pyridine compound (1) is
15 then attached to a trityl chloride resin solid support, to form compound (2).

Before, after, or simultaneously with the formation of compound (2), a second molecule of 3-substituted pyridine
20 compound (1) is reacted firstly with mesyl chloride, and secondly with m-chloroperbenzoic acid to form compound (3) of formula $\text{O}^-\text{NC}_5\text{R}_4\text{-R}'\text{-O-Ms}$, wherein Ms is a mesyl group.

Next, compound (2) is reacted with one equivalent of compound
25 (3) to form 1,3-alkylpyridinium compound (4a) having the formula $\text{O}^-\text{NC}_5\text{R}_4\text{-R}'\text{-I}^+\text{NC}_5\text{R}_4\text{-R}'\text{-O-SUPPORT}$, i.e. two pyridine-containing monomer units are combined to form a dimeric 1,3-alkylpyridinium compound, the -Ms leaving group of compound (3) reacting with the nitrogen atom of the pyridine group of
30 compound (2). This step is repeated the appropriate number of times to obtain an 1,3-alkylpyridinium compound end product having the desired degree of polymerisation, i.e. compound (2) is reacted with 1,3-alkylpyridinium compound (4a) to produce

compound (4b) of formula $\text{O}^-\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{I}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{O-SUPPORT}$, wherein n is preferably 20 to 100.


As discussed above, in preferred embodiments of the present invention, chain extension to produce oligomers having a higher degree of polymerisation may be accelerated by releasing oligomers having a lower degree of polymerisation from the solid support as compounds having the formula (5), and reintroducing them as reagents as an alternative to, or in addition to, the second pyridine compound used in step (b). Thus, in these preferred embodiments, a compound of formula $\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{-X}$ (5) may be converted to a compound of formula $\text{O}^-\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{-Z}$ (5a) per step (b), and the compound of formula (5a) may then be reacted with the compound of formula (2) formed in step (a) or (c), per step (d).

Finally in Scheme 1, compound (4b) is detached from the solid support, and reduced to form 1,3-alkylpyridinium compound (5) of formula $\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Cl}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{-OH}$ using hydrochloric acid, as shown. The counter ion in final product (5) is thus chloride.

As referred to above, the method of the present invention is advantageous in that a variety of linker groups may be used, which may incorporate different functionalities.

Reaction Scheme 2 below describes a method for producing a compound from which starting material $\text{NC}_5\text{R}_4\text{-R}'\text{-X}$ used in step (a) of the present invention may in turn be produced, in which R' is an alkenyl-containing linking group having either cis- or trans- isomerism.

Scheme 2




wherein: TBDMS=t-butyldimethylsilyl
 DMAP=4-dimethylaminopyridine
 THF=tetrahydrofuran
 LDA=lithium diisopropylamine
 TBAF=t-butylammonium fluoride

15

Reaction Scheme 3 below describes a method for producing a compound from which starting material $\text{NC}_5\text{R}_4\text{-R}'\text{-X}$ of formula 20 (1) used in step (a) of the present invention may in turn be produced, in which R' is an alkylene linking group having a pendant alcohol group to which a fluorescent group may be attached.

25

Scheme 3



wherein: Ph=phenyl
Py=pyridine
Tr=trityl
Bn=benzyl

5

As shown in Scheme 3, triol (6) is differentially protected to form alcohol (7), which is subsequently converted to bromide (8), which can then in turn be used to form starting material $\text{NC}_5\text{R}_4\text{-R}'\text{-X}$ of formula (1) used in step (a) of the
10 present invention.

Fluorescent labels, available commercially, can be reacted with the monomer or oligomer, depending on the reaction conditions necessary. To avoid cell damage, the excitation
15 wavelength of the fluorescent label is preferably above 350 nm, and the emission wavelength should preferably be greater than 80 nm higher, to permit ratiometric measurements. Suitable fluorescent labels include fluorescein, lissamine, and rhodamine based compounds, although other fluorescent
20 materials may be used according to conditions. Such a linking group having a fluorescent group would preferably be incorporated into the 1,3-APS compound end product only once or twice.

CLAIMS

1. A method of producing a linear di-substituted pyridinium compound, the method comprising the steps of:

5 (a) attaching a first 2, 3, or 4-substituted pyridine compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-X}$ (1) to a solid support, to form a compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-Y-SUPPORT}$ (2), wherein SUPPORT represents the solid support, R is selected from hydrogen, hydroxyl, and substituted or unsubstituted alkyl, alkoxy, aryl, alkaryl, aralkyl, and alkenyl groups, R' is a
10 first linking group, X is a group which can react with the solid support to attach the first pyridine compound to the support, and Y is absent or is a second linking group;

(b) forming a di-substituted pyridine compound of the
15 formula $\text{A-}^+\text{NC}_5\text{R}_4\text{-R}'\text{-Z}$ (3) from a second 2, 3, or 4-substituted pyridine compound of formula (1), which second pyridine compound may be the same as or different to the first pyridine compound, wherein A is a protecting group, and Z is a leaving group;

20 (c) reacting the compound of formula (2) formed in step (a) with the compound of formula (3) formed in step (b), to form a di-substituted pyridinium compound of the formula
 $\text{A-}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^-\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{Y-SUPPORT}$ (4), wherein Q⁻ is a counter ion and n=1;

25 (d) optionally, repeating step (c) as many times as required to obtain a compound of formula (4) wherein n is an integer of 2 or greater; and

(e) detaching the compound of formula (4) from the solid support, and reducing to form a di-substituted pyridinium
30 compound of the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^-\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{-X}$ (5), wherein n is an integer, and Q⁻ and X are a counter ion and a group which can react with the solid support to attach the first pyridine compound to the support respectively, which may the

same or different as Q and X defined above in steps (c) and (a) respectively.

2. A method according to claim 1 wherein each R group is 5 hydrogen.

3. A method according to claim 1 or 2 wherein in step (d), step (c) is repeated such that in formula (5) $n=20$ to 100.

10 4. A method according to claim 1, 2 or 3 wherein the compound of formula (1) is prepared by reaction of a compound of formula $Z'-R''-X$ with a pyridine compound, with protection of the X- group as necessary, wherein R'' is a linker group and Z' is a suitable leaving group.

15

5. A method according to claim 4 wherein the compound of formula (1) is prepared by reacting $Br-R''-OH$ with t-butyldimethyl-chlorosilane (TBDMSCl) to form $Br-R''-OTBDMS$, which is reacted with 2,3 or 4-methylpyridine (2,3, or 4-
20 picoline) with deprotection of the X group to form $NC_5R_4-R'-OH$.

6. A method according to any one of claims 1 to 5 wherein linker group R' has no terminal carbon atoms.

25

7. A method according to any preceding claim wherein each group R' is the same or different and is selected from an alkylene group, an alkenyl-containing group, and a cyclopropanyl-containing group.

30

8. A method according to claim 7 wherein each group R' is the same or different and is selected from a group $-(CH_2)_m-$, wherein m is an integer from 2 to 12, a group having from 2

to 12 carbon atoms containing one or more alkenyl groups, and
cis- or trans- $-(CH_2)_p$ -cyclopropanyl- $(CH_2)_q$ - wherein p and q
are the same or different and are integers from 1 to 4.

5 9. A method according to any preceding claim wherein R' is
a group which comprises a fluorescent group, or a group to
which a fluorescent group can be attached.

10. A method according to claim 9 wherein R' has a pendant
10 alcohol group, for attachment of a fluorescent group.

11. A method according to any preceding claim wherein in
formula (1) X is selected from hydroxyl, carboxyl, thiol, and
amine groups.

15

12. A method according to claim 11 wherein X is a hydroxyl
group.

13. A method according to claim 12 wherein the compound of
20 formula (1) is a compound of the formula $NC_5R_4-(CH_2)_n-OH$.

14. A method according to any preceding claim wherein the
solid support material comprises an organic resin having
functionality which can react with group X of the compound of
25 formula (1).

15. A method according to claim 14 wherein the solid support
material comprises trityl chloride or a functionalised
polystyrene resin.

30

16. A method according to any preceding claim wherein group
Y in formula (2) is an oxygen atom.

17. A method according to any preceding claim wherein in step (b) group X is converted to a mesyl (methanesulphonyl) group by reaction with mesyl chloride.

5 18. A method according to any preceding claim wherein A is oxygen or BH_3^- .

19. A method according to claim 18 wherein A is oxygen, and the nitrogen atom of the second pyridine compound of formula
10 (1) used in step (b) is converted to the N-oxide by reaction of the nitrogen atom of the pyridine group with a peracid.

20. A method according to claim 19 wherein the peracid comprises m-chloroperbenzoic acid.

15

21. A method according to any preceding claim wherein counter ion Q^- in step (c) is an iodide ion.

22. A method according to any preceding claim wherein
20 oligomers having the formula (5) are released from the solid support and reintroduced as reagents as an alternative to, or in addition to, the second pyridine compound used in step (b).

23. A method according to claim 22 wherein a compound of
25 formula $\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}']_n\text{-X}$ (5) is converted to a compound of formula $\text{A-}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}']_n\text{-Z}$ (5a) per step (b), and the compound of formula (5a) is then reacted with the compound of formula (2) formed in step (a) or (c), per step (d).

30

24. A method according to any preceding claim wherein the compound of formula (4) is detached from the solid support, and reduced to form a di-substituted pyridinium compound of

the formula $\text{NC}_5\text{R}_4\text{-R}'\text{-}[\text{Q}^+\text{NC}_5\text{R}_4\text{-R}'\text{-}]_n\text{-X}$ (5) using an acid.

25. A method according to claim 24 wherein the acid is hydrochloric acid, and counter ion Q^- is chloride.

5

26. A method according to any preceding claim wherein the di-substituted di-substituted pyridinium compound is a linked dialkyl pyridinium compound.

10 26. A method of producing a linear di-substituted pyridinium compound substantially as hereinbefore described with reference to Reaction Schedule 1.

PCT/GB2004/002666



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.